

The potential of typhoid conjugate vaccines in Burkina Faso

Typhoid, a serious enteric fever spread through contaminated food and water, is a substantial public health issue that disproportionately impacts children and marginalized populations in Asia and sub-Saharan Africa. The Global Burden of Disease (GBD) study estimates that, in 2017, there were nearly 11 million typhoid cases and more than 116,000 typhoid deaths worldwide.¹ Additionally, strains of drug-resistant typhoid are spreading, causing global concern.²

TYPHOID CONJUGATE VACCINES


Typhoid vaccination can reduce the need for antibiotics, slow expansion of drug-resistant strains, and save lives. Newly licensed and World Health Organization (WHO)-prequalified typhoid conjugate vaccines (TCVs) have several advantages over earlier typhoid vaccines. They:

- provide longer-lasting protection;
- require only one dose; and
- are suitable for young children over 6 months.

These qualities will allow better protection for younger children and expanded coverage through routine childhood immunization programs.

WHO RECOMMENDATION AND GAVI SUPPORT

In March 2018, WHO recommended that typhoid-endemic countries introduce prequalified TCVs into routine childhood immunization programs as a single dose for infants and children over 6 months of age, accompanied by catch-up vaccination campaigns for children up to 15 years of age, where feasible. Additionally, WHO recommended prioritizing countries with a high burden of disease and/or a growing burden of drug-resistant typhoid and in response to confirmed typhoid outbreaks. Gavi, the Vaccine Alliance has earmarked US\$85 million to support the introduction of TCVs into routine immunization programs and is accepting applications for financial support, with introductions anticipated as soon as 2019.



According to GBD estimates, Burkina Faso had 431 typhoid cases per 100,000 population in 2017—the highest typhoid incidence rate in Africa and the second highest in the world.

AN OPPORTUNITY FOR BURKINA FASO

TCVs could have a substantial benefit in Burkina Faso, where typhoid inflicts a significant health and economic burden. The GBD estimates that, in 2017, Burkina Faso had:

- **90,931 typhoid cases** or **431 cases per 100,000 population**, 64 percent of which were among children under 15 years of age; and
- **1,150 typhoid deaths**, 69 percent of which were among children under 15 years of age.¹

A recent modeling study⁴ shows that a catch-up campaign up to 15 years of age followed by routine immunization is the preferred strategy and likely to be cost-effective in Burkina Faso.*

*At a willingness-to-pay threshold of \$100 or more to avert one disability-adjusted life-year (DALY).



Bill & Melinda Gates Foundation/Sam Reinders

Typhar-TCV® typhoid conjugate vaccine was prequalified by the World Health Organization in December 2017.



Sabin Vaccine Institute

Four-year-old Golden Kondowe was the first child in Africa to receive a vaccine as part of the TyVAC effectiveness study in Malawi, February 2018.

TyVAC STUDY IN BURKINA FASO

In order to build evidence of the effectiveness of TCVs in protecting children from typhoid, researchers with the Typhoid Vaccine Acceleration Consortium (TyVAC) are conducting four different studies in Bangladesh, Burkina Faso, Malawi, and Nepal. In Burkina Faso, TyVAC and project partners are planning to study how well TCVs produce an immune response to typhoid in children between 9 months and 2 years of age as well as the safety of the vaccine when given alone or alongside other routine childhood vaccines. While WHO already recommends TCV introduction in all typhoid-endemic countries, this additional evidence will help inform ongoing decisions about TCV vaccination in low- and middle-income countries.

REFERENCES

1. Global Burden of Disease. *The Lancet*. 2018.
2. Wong VK, Baker S, Pickard DJ, et al. Phylogeographical analysis of the dominant multidrug-resistant H58 clade of *Salmonella* Typhi identifies inter- and intracontinental transmission events. *Nature Genetics*. 2015;47(6):632-639.
3. Riewpaiboon A, Piatti M, Ley B, et al. Cost of illness due to typhoid fever in Pemba, Zanzibar, East Africa. *Journal of Health, Population and Nutrition*. 2014;32(3):377-385.
4. Bilcke J et al. Cost-effectiveness of routine and campaign use of typhoid Vi-conjugate vaccine in Gavi-eligible countries: a modelling study. *The Lancet Infectious Diseases*. Accepted.

Learn more and join the effort at www.takeontyphoid.org.

#TakeOnTyphoid